

Eclipsing Binaries With Possible Light-Time Effect *

Petr Zasche

Astronomical Institute, Charles University, Prague, Czech Republic

Miloslav Zejda

Nicolas Copernicus Observatory and Planetarium, Brno, Czech Republic

Luboš Brát

Velká Úpa 193, Pec pod Sněžkou, Czech Republic

Abstract. The period changes of six eclipsing binaries have been studied with focus on the Light-time effect. With the least squares method we also calculated parameters of such effect and properties of the unresolved body in these systems. With these results we discussed probability of presence of such bodies in the systems with respect to possible confirmation by another method. In two systems we also suggested a hypothesis of fourth body or magnetic activity for explanation of the "second-order variability" after subtraction of the light-time effect of the third body.

Keywords: stars: binaries: eclipsing – stars: individual – multiple stars – period variations

1. Introduction

Eclipsing binary systems provide a good opportunity for studying the presence of an unresolved third body by observing their minima times because of the light-time effect (hereafter LITE). It was explained by Irwin (1959) and its necessary criteria have been mentioned by Frieboes-Conde & Hertzeg (1973) and also by Mayer (1990). Presence of the third body in the system is possible only if the times of minima behave in agreement with a theoretical LITE curve, the resultant mass function has reasonable value and corresponding variations in radial velocities are measured. In the last decade also a confirmation by astrometry seems to be plausible.

In each case we have calculated new light elements of the eclipsing pair and also the parameters of the predicted third body orbit. The tables I. and II. present results for each system, where M_i are masses of components, p_i computed period of the unresolved body, A_i semi-amplitude of LITE, e_i eccentricity, ω_i length of periastron, $f(M_i)$ mass function and $M_{i,min}$ minimum mass (for $i = 90^\circ$) of predicted body, respectively. The subscript 3 and 4 denotes the parameter of the third and fourth body, respectively.

* For further details, please contact PZ, e-mail address: petr.zasche@email.cz






Figure 1. The O-C diagrams of selected eclipsing binaries. The curves represent proposed light-time effect caused by the third and fourth bodies with periods and semiamplitudes given in the tables I and II. The individual primary and secondary minima are denoted by dots and circles, respectively. Larger symbols correspond to the photoelectric or CCD measurements which were given higher weights in our computations. In OO Aql and T LMi cases the LITE caused by the third body is represented by the dashdotted line together with the LITE caused by the fourth body by the solid line. The residuals after subtraction of LITE caused by the third body are shown in figures on the right.

2. Investigated systems

We have derived new linear light elements:

OO Aql: Min I = HJD 24 39322.6902 + 0^d.50679191 · E,

V338 Her: Min I = HJD 24 41945.3324 + 1^d.30574608 · E,

T LMi: Min I = HJD 24 23856.4200 + 3^d.01988799 · E,

RV Lyr: Min I = HJD 24 45526.4003 + 3^d.59901148 · E,

TW Lac: Min I = HJD 24 24387.2547 + 3^d.03749292 · E,

V396 Mon: Min I = HJD 24 34769.4234 + 0^d.39634326 · E.

In the case of OO Aql and T LMi also the hypothesis of the fourth body was suggested. The results are in Table II. In the system OO Aql the components are very similar and also similar to the Sun (sp. G5V, $T \simeq 5700$ K), so the second order variations could also be caused by the the magnetic cycles, similar to the Sun (so-called Applegate mechanism, see eg. Applegate (1992)).

In the system V338 Her also the quadratic term was applied, so the mass transfer in the system could be present. From this hypothesis (with mass transfer parameter $q = 2 \cdot 10^{-10}$) we have derived the mass transfer rate to $\simeq 4 \cdot 10^{-8} M_{\odot}/\text{yr}$.

In a few cases here the potential third body would be detectable in the detailed light curve analysis. Especially the cases where the third body has not negligible mass, comparing with the eclipsing pair. Regrettably we have no information about the distance of individual systems and also about the absolute magnitudes, so the determination of the angular separation of the possible third body is very uncertain.

Table I. The results for six investigated LITE systems.

Name of star	Spectrum	$M_1 + M_2$ [$M_{\odot}$]	p_3 [years]	A_3 [days]	e_3	ω_3 [deg]	$f(M_3)$ [$M_{\odot}$]	$M_{3,min}$ [$M_{\odot}$]	Ref.
OO Aql	G5V+G5?	1.05+0.88	71.72	0.0369	0.13	0.0	0.052	0.714	[1]
V338 Her	F2V+K0	1.42+0.40	26.35	0.0150	0.59	128.7	0.031	0.562	[2]
T LMi	A0+G5III	6.10+0.60	136.98	0.1023	0.29	24.6	0.329	3.177	[3]
RV Lyr	A5+K4III	3.70+1.30	67.99	0.0331	0.31	138.5	0.044	1.192	[4]
TW Lac	A2IV	2.80+1.87	105.98	0.0844	0.70	213.1	0.523	3.184	[5],[2]
V396 Mon	?F8+K8?	1.20+0.47	133.71	0.0519	0.41	114.2	0.042	0.603	[6]

References: [1] – Al-Naimiy et.al. (2000); [2] – Budding (2004); [3] – Cester (1979); [4] – Budding (1984); [5] – Halbedel (1984); [6] – Yang & Liu (2001).

Table II. Parameters for the proposed fourth components.

Name of star	p_4 [years]	A_4 [days]	e_4	ω_4 [deg]	$f(M_4)$ [$M_\odot$]	$M_{4,min}$ [$M_\odot$]
OO Aql	19.71	0.0034	0.01	31.7	0.001	0.159
T LMi	37.42	0.0144	0.17	44.8	0.011	1.109

3. Conclusion

We have derived new LITE parameters for six eclipsing binaries by means of an $O-C$ diagram analysis. In two cases, OO Aql and T LMi, another variation was found, so there is a possibility of a presence of the fourth body in the system, or magnetic activity in them. But we have not enough data to make a final decision. So the consequence is, that for the confirmation of presence of the LITE in these systems, we need detailed photometric, spectroscopic or astrometric data of these binaries.

4. Acknowledgements

This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France, and of NASA's Astrophysics Data System Bibliographic Services. This investigation was supported by the Czech-Greek project of collaboration RC-3-18 of Ministry of Education, Youth and Sport and by the Grant Agency of the Czech Republic, grant No. 205/04/2063

References

- Applegate, J.H.: 1992, *AJ* **385**, 621.
 Budding, E.: 1984, *BICDS* **27**, 91.
 Budding, E.: 2004, *A&A* **417**, 263.
 Cester, B. et.al.: 1979, *A&A* **36**, 273.
 Frieboes-Conde, H.; Hertzeg, T.: 1973, *A&AS* **12**, 1.
 Halbedel, E.M.: 1984, *IBVS* **2549**, 1.
 Irwin, J.B.: 1959, *AJ* **64**, 149.
 Mayer, P.: 1990, *BAICz* **41**, 231.
 Naimiy, H.M.K.; Al-Masharfeh, T.H.S.: 2000, *ApSS* **273**, 83.
 Yang, Y.; Liu, Q.: 2001, *AJ* **122**, 425.